



Research Article

Metabolic Health Outcomes Following Nine Months of Mild Caloric Restriction in Male Rats Adhering To a Western or Vegan Diet

Richard J Bloomer*, John Henry M Schriefer, Trint A Gunnels and Matthew Butawan

School of Health Studies, University of Memphis, Memphis, TN, USA

Abstract

Background: Both diet composition and total calorie intake can influence body mass and measures of metabolic health.

Methods: We assigned male Long-Evans rats (N=28) to either a high-fat Western Diet (WD) or a moderate fat, purified Vegan Diet (VD) for 12 months, the first 3 of which food was provided ad libitum and the latter 9 in which food was restricted to approximately 90% of animals' daily needs. Half of the animals in each diet group were assigned to be exercise trained (+E) by running on a treadmill three days per week and the other half served as sedentary controls. Body mass was monitored daily and reported at three month intervals; plasma indicators of metabolic health were determined at the conclusion of the 12-month period.

Results: After three months of ad libitum feeding, body mass was higher for the WD compared to the VD (549g vs 484g; $p < 0.00001$; pooled data for both exercise groups), and slightly less for the exercise-trained animals as compared to the sedentary animals (494g vs 539g; $p = 0.006$; pooled data for both diet groups). Following 9 months of mild caloric restriction, body mass decreased approximately 15% for all groups, except for the WD+E group, for which the decrease was approximately 8%. Oxidative stress biomarkers, as measured by advanced oxidation protein products ($p < 0.0001$) and malondialdehyde ($p = 0.05$), were lower for the VD groups as

*Corresponding author: Richard J Bloomer, School of Health Studies, 106 Roane Fieldhouse, University of Memphis, Memphis, TN 38152, USA, Tel: +1 9016785638; Fax: +1 9016783591; E-mail: rbloomer@memphis.edu

Citation: Bloomer RJ, Schriefer JHM, Gunnels TA, Butawan M (2020) Metabolic Health Outcomes Following Nine Months of Mild Caloric Restriction in Male Rats Adhering To a Western or Vegan Diet. J Altern Complement Integr Med 6: 097.

Received: April 02, 2020; **Accepted:** April 06, 2020; **Published:** April 13, 2020

Copyright: © 2020 Bloomer RJ, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

compared to the WD groups. Triglycerides ($p < 0.0001$) and cholesterol ($p = 0.0006$) were also lower for the VD groups as compared to the WD groups, but glucose was not ($p = 0.09$).

Conclusion: While mild caloric restriction results in a similar percent decrease in body weight in animals adhering to both a WD and VD, the improved metabolic health of animals following the VD, as indicated by lower plasma biomarkers of oxidative stress and blood lipids, highlights the fact that consuming a purified diet yields benefits that extend beyond mere weight loss.

Keywords: Caloric restriction; Cholesterol; Dietary restriction; Fasting; Oxidative stress; Triglycerides; Vegan body mass

Introduction

Aside from caloric restriction, manipulating diet composition by adjusting macronutrient ratios and types can have a profound influence on overall health outcomes [1-4]. One form of manipulation involves veganism. This pattern of eating eliminates all animal products and has been reported to yield favorable outcomes related to cardio-metabolic health, including reductions in body mass [5-7]. Unlike caloric restriction, which typically calls for a significant reduction in daily dietary energy needs, veganism places no limitation on calories and allows for ad libitum food intake.

We have studied a very stringent form of vegan-based dieting over the past several years, with reductions of body mass, blood lipids and measures of oxidative stress in as little as three weeks [8-12]. This approach, referred to as the "Daniel Fast" plan, allows for ad libitum food intake but places firm restrictions on the type of food that is allowed, with choices primarily limited to fruits, vegetables, whole grains, legumes, nuts, seeds and plant-based oils. No alcohol, sweeteners or refined foods are allowed resulting in carbohydrate sources that are complex with low glycemic indices. By default, this plan has an abundance of dietary fiber and plant-derived fatty acids, and relatively high concentrations of antioxidants.

While altering the diet composition in a way similar to that described above can be useful as a health promotion strategy, the most commonly utilized approach is to simply reduce calorie intake the classic "hypocaloric" dieting. While reducing calories may be achieved using a variety of different diet plans (e.g., simple caloric restriction, intermittent or alternate day fasting), the majority of plans do in fact result in a lower total calorie intake, leading to a reduction in body weight. However, while weight loss is often achieved by many dieters, what is often not considered in particular, outside of the context of a controlled research study or a clinical case is how diet composition may impact other measures of health; that is, those that may not be "visible" through a simple assessment of body mass.

The present study sought to determine the influence of mild, chronic caloric restriction in male rats. Outcome variables included body mass but also biochemical measures of health that are important

but often not considered by those adopting a weight loss diet. We hypothesized that intake of a purified diet would yield results that were more favorable as compared to a Western Diet, in particular with regards to the biochemical outcomes. Moreover, we included thrice weekly treadmill walking/running in the model to determine the potential additive influence of exercise on the outcome measures.

Methods

Animals and housing

Male Long-Evans rats (N=28) were purchased from Harlan Laboratories, Inc. (Indianapolis, IN) at the age of 3-4 weeks. Rats were individually housed in a climate controlled room (21°C) with a 12:12-h light-dark cycle. They were initially fed a standard rat chow (Harlan 1018) with ad libitum water, and then transitioned to the assigned diet after two weeks. During this period the rats were familiarized with the treadmill on three separate days. All experimental procedures were approved by The University of Memphis *Institutional Animal Care and Use Committee*.

Diet group assignment

The rats were randomly assigned to one of four intervention groups: Western Diet with exercise (WD+E; n=7); Western Diet without exercise (WD; n=7); Vegan Diet with exercise (VD+E; n=7); Vegan Diet without exercise (VD; n=7). Both diets (provided in pellet form) were purchased from Research Diets, Inc. (New Brunswick, NJ). The WD (D12079B) was formulated to mimic a typical human WD, containing 17% protein, 43% carbohydrates (sucrose and corn) and 40% fat (butter and corn oil), whereas the VD included 15% protein, 60% carbohydrates, and 25% fat. The nutrient composition of both diets has been presented previously [13]. The types and quantities of macronutrient sources in the VD were based on the diet plans of the human subjects in our prior studies using a vegan diet [8,10,11]. The VD was rich in inulin+cellulose and flax oil, contributing to the carbohydrate and lipid content, respectively. The dietary intervention period was 12 months in duration. During months 1-3, food was provided ad libitum, while during months 4-12, the rats were weighed daily and provided a quantity of food thought to maintain them at approximately 90% of their free-feeding body mass measured just prior to food restriction. Water was provided ad libitum throughout the 12-month period.

Treadmill exercise

Animals in the exercise groups performed walking/running exercise on a level motorized treadmill three days per week, with a progressive increase in speed and duration. Specifically, the animals began training at 20 m·min⁻¹ for 15 min·day⁻¹ (week 1), progressed to 25 m·min⁻¹ for 30 min·day⁻¹ (week 2) and 25 m·min⁻¹ for 35 min·day⁻¹ (weeks 3-52).

Outcome measures

Body mass was measured daily using a Mettler Toledo PG2002-S balance equipped with dynamic weighing. At the end of the 12-month intervention, rats were euthanized via CO₂ inhalation and blood was collected from the inferior vena cava into vacutainer tubes containing EDTA. Plasma was separated and aliquots stored at -70°C for analysis of metabolic health biomarkers. Advanced Oxidation Protein Products (AOPP) was measured using the methods described by

the reagent manufacturer (Cell Biolabs, Inc. San Diego, CA; product #: STA-318). Malondialdehyde (MDA) was analyzed following the procedures of Jentzsch et al., (1996) using reagents purchased from Northwest Life Science Specialties (Vancouver, WA; product #: NWK-MDA01). Triglycerides (TAG), total cholesterol, and glucose were analyzed following standard enzymatic procedures as described by the reagent manufacturer (Thermo Electron Clinical Chemistry). All samples were analyzed in duplicate.

Statistical analysis

All plasma variables were analyzed using a one-way Analysis of Variance (ANOVA). Contrasts were used to compare individual groups with regards to bloodborne variables and body mass. All analyses were performed using JMP statistical software (SAS Institute; Cary, NC). Statistical significance was set at p ≤ 0.05. All data are expressed as the mean ± SD.

Results

Following the initial three months of ad libitum feeding, body mass was higher for animals in the WD groups as compared to the VD groups, and slightly less for the exercise-trained animals as compared to the sedentary animals. Following the 9-month period of mild caloric restriction, body mass decreased in all groups, with an approximate 15% decrease for the WD, VD and VD+E groups, and an approximate 8% decrease for the WD+E group (Table 1). After the 12-month intervention, pooled data for both VD groups were lower compared to WD groups for advanced oxidation protein products (p<0.0001) and malondialdehyde (p=0.05), and for triglycerides (p<0.0001) and cholesterol (p=0.0006), with a trend for lower glucose (p=0.09). Biochemical data for all groups and the noted differences are displayed in table 2.

	Western Diet + Exercise	Western Diet	Vegan Diet + Exercise	Vegan Diet
Month 0 (Baseline)	187±9	187±12	193±8	185±14
Month 3	519±28	579±37	471±47	498±36
Month 6	487±28	515±22	425±26	432±22
Month 9	478±23	516±27	414±26	403±76
Month 12	475±24*†	490±22*§	398±27*	412±19*

Table 1: Body mass (g) of male rats assigned to two different diets with and without exercise for 12 months.

Values are mean ± SD.

* Significant difference between months 3 and 12 for WD+E (p=0.03), WD (p<0.0001), VD+E (p=0.0001) and VD (p<0.0001)

† Significant difference at 12 months between WD+E and VD+E (p=0.0002); WD+E and VD (p=0.002)

§ Significant difference at 12 months between WD and VD+E (p<0.0001); WD and VD (p<0.0001)

Discussion

To our knowledge, this is the first study to determine the combined influences of dietary composition and exercise during a period of caloric restriction on outcomes specific to metabolic health. There are several key findings from this study. Consumption of an ad libitum WD leads to greater weight gain as compared to a VD. Exercise reduces body mass gain during the ad libitum period, and particularly for the WD. Despite a similar percentage weight loss for the WD

and VD groups during the period of caloric restriction, plasma lipids and indicators of oxidative stress were significantly lower for the VD. During the period of caloric restriction, exercise had little impact on plasma markers of oxidative stress and lipids, with the exception of AOPP and triglyceride levels with the WD.

	Western Diet + Exercise	Western Diet	Vegan Diet + Exercise	Vegan Diet
AOPP ($\mu\text{mol}\cdot\text{L}^{-1}$)	114±73	151±71	22±14	22±11
Malondialdehyde ($\mu\text{mol}\cdot\text{L}^{-1}$)	4.0±2.0	3.5±2.6	1.8±0.7	1.8±0.1
Triglyceride ($\text{mg}\cdot\text{dL}^{-1}$)	90±44	151±62	32±9	37±19
Cholesterol ($\text{mg}\cdot\text{dL}^{-1}$)	94±14	90±21	63±8	68±12
Glucose ($\text{mg}\cdot\text{dL}^{-1}$)	121±34	127±22	97±17	107±17

Table 2: Biochemical data of male rats assigned to two different diets with and without exercise for 12 months.

Values are mean ± SD.

AOPP (Advanced Oxidation Protein Products): Significant difference between WD+E and VD+E ($p=0.002$) and VD ($p=0.002$); between WD and VD+E ($p<0.0001$) and VD ($p<0.001$)

Malondialdehyde: Significant difference between WD+E and VD+E ($p=0.03$) and VD ($p=0.03$)

Triglyceride: Significant difference between WD+E and VD+E ($p=0.01$) and VD ($p=0.02$); between WD and VD+E ($p<0.0001$) and VD ($p<0.001$)

Cholesterol: Significant difference between WD+E and VD+E ($p=0.0005$) and VD ($p=0.002$); between WD and VD+E ($p=0.001$) and VD ($p=0.006$)

The lower weight gains during the period of ad libitum feeding support observations that vegetarian diets are associated with better weight maintenance [14]. Although the amount of food consumed by each animal was not measured during the period of ad libitum feeding, we believe that the volume was similar, as rats have been reported to consume relatively equal amounts of food, despite differences in caloric density [15].

Interestingly, regardless of the type of food consumed, when animals were subjected to a mild caloric restriction, all experienced a decrease in body mass of approximately 15%, except for the WD+E group, for which the reduction was approximately 8%. The lower percent loss could have been due to the fact that the exercise in the WD group attenuated the weight gain (i.e., weight gain in the WD group without exercise was far greater) and there was less weight to lose in the WD+E group. From a practical perspective, it appears that regardless of food composition, body mass can be reduced following a period of mild caloric restriction. This should be viewed as encouraging, in particular for those individuals who may not have the desire to alter what they eat but may be capable of reducing the quantity of food consumed.

As alluded to above, engaging in thrice weekly aerobic exercise resulted in a reduction in the amount of body weight gained during the initial three months of ad libitum feeding, which was expected based on prior knowledge of exercise as a therapeutic tool [16]. We anticipated that the exercise groups would have a lower overall body mass due to the increased caloric expenditure from the exercise bouts. For example, animals that did not exercise and were assigned to the WD gained 18% more weight, while those who did not exercise and were assigned to the VD gained 13% more weight. Exercise results in energy expenditure and is a commonly used modality to aid overall cardio-metabolic health [17]. Multiple studies support the role of

exercise as an adjunct to diet therapy for purposes of weight management [18]. Interestingly and as shown in table 1, similar findings have been noted in a recent animal study, in which rats placed on an 8% caloric restriction, high fat diet experienced similar weight reduction with or without exercise [19].

As mentioned above, the percentage weight loss for the WD and VD groups was similar during the restriction period. However, despite this, oxidative stress biomarkers and blood lipids were significantly lower in animals adhering to the VD. Often, individuals engage in a dietary program with the goal of losing weight and becoming “healthier”. In a non-clinical/research setting, the former is easy to determine; individuals can simply stand on a scale. The latter is much more complex but often is determined by the lay person simply based on how they feel. Our data show that despite a similar degree of weight loss during the nine-month restriction period, outcomes related to metabolic health are very different depending on the type of food that is consumed.

For example, protein oxidation, as measured by AOPP, was approximately 6-7 folds higher for the WD groups as compared to the VD groups, while MDA values were approximately 2 fold higher. Findings of lower oxidative stress in vegetarians have been noted previously [20]. Oxidative stress is associated with cardiovascular and neurodegenerative disease, as well as most other human diseases and minimizing the overall oxidative burden is generally viewed as a favorable health outcome [21,22]. Due to the fact that body mass (obesity) has been associated with oxidative stress, it is possible that the elevated values in relation to animals consuming the WD are a function of their higher absolute body mass [23]. That is, despite the fact that the WD and VD animals experienced a similar overall percent decrease in body mass during the restriction period, the absolute body mass of the WD animals at the conclusion of the 12-month intervention remained approximately 20% higher as compared to the VD.

While this difference in body mass may have contributed to our findings, an alternative hypothesis is that the diet composition was responsible for the differing oxidative stress values. Indeed, high intake of saturated fat and simple sugar as found in the WD, is associated with increased production of reactive oxygen species and related negative outcomes [24,25]. Moreover, flax oil and dietary fiber two components of the VD have been reported to provide favorable effects with regards to oxidative stress [26,27]. In the same way as for oxidative stress variables, both triglycerides and cholesterol were lower for the VD following the restriction period, with non-statistically significant lowering for glucose. These results are logical, as reducing intake of saturated fat, cholesterol and simple sugar is often met with a lowering in blood triglycerides and cholesterol.

Finally, as with the percentage of weight lost during the restriction period, exercise had little impact on oxidative stress and lipid levels; the exception being a lowering of triglyceride levels when adhering to the WD. Physical activity has been reported to result in lower levels of oxidative stress, with mixed findings for blood lipids [28]. It is possible that the impact of the diet restriction overwhelmed any effect of the exercise, to the extent that it was not observed in our sample of animals. Alternatively, the exercise in the present study may have been of too low of an intensity or volume to alter oxidative stress or cholesterol. Future work is needed to replicate our study and determine the independent and combined role of aerobic exercise, perhaps of higher volume and intensity, on measures of oxidative stress and

blood lipids when animals/individuals adhere to a Western diet or purified vegan diet.

Conclusion

To our knowledge, this is the first study to investigate the impact of both dietary restriction and exercise, inclusive of ad libitum feeding and caloric restriction, on measures of metabolic health. The findings suggest that macronutrient composition, and not simply caloric intake, has an influence on the degree of oxidative stress, as well as blood lipids. These findings underscore the value of consuming a purified diet if the main objective is not simply to lose weight but to become healthy overall from a metabolic perspective.

Conflict of Interest

The authors declare no conflicts of interest related to this work.

Author Contributions

RJB was responsible for the study design, biochemical analyses, statistical analyses and manuscript preparation. JMS and TAG were responsible for animal training, data collection, database management, and assistance with the study design and manuscript editing. MB was responsible for manuscript preparation. All authors read and approved of the manuscript.

Acknowledgement

Funding for this work was provided by the University of Memphis. Appreciation is extended to Drs. Sang-Rok Lee, Randal Budington, and Karyl Buddington for assistance in data collection.

References

1. Das SK, Balasubramanian P, Weerasekara YK (2017) Nutrition modulation of human aging: the calorie restriction paradigm. *Mol Cell Endocrinol* 455: 148-157.
2. Golbidi S, Daiber A, Korac B, Li H, Essop MF (2017) Health benefits of fasting and caloric restriction. *Curr Diab Rep* 17: 123.
3. Makki K, Deehan EC, Walter J, Bäckhed F (2018) The impact of dietary fiber on gut microbiota in host health and disease. *Cell Host Microbe* 23: 705-715.
4. Shondelmyer K, Knight R, Sanivarapu A, Ogino S, Vanamala JKP (2018) Focus: Nutrition and Food Science: Ancient Thali Diet: Gut Microbiota, Immunity, and Health. *Yale J Biol Med* 91: 177.
5. Glick-Bauer M, Yeh MC (2014). The health advantage of a vegan diet: exploring the gut microbiota connection. *Nutrients* 6: 4822-4838.
6. Appleby PN, Key TJ (2016) The long-term health of vegetarians and vegans. *Proc Nutr Soc* 75: 287-293.
7. Kahleova H, Levin S, Barnard N (2017) Cardio-metabolic benefits of plant-based diets. *Nutrients* 9: 848.
8. Bloomer RJ, Kabir MM, Canale RE, Trepanowski JF, Marshall KE, et al. (2010) Effect of a 21 day Daniel Fast on metabolic and cardiovascular disease risk factors in men and women. *Lipids Health Dis* 9: 94.
9. Trepanowski JF, Kabir MM, Alleman RJ Jr, Bloomer RJ (2012) A 21-day Daniel fast with or without krill oil supplementation improves anthropometric parameters and the cardiometabolic profile in men and women. *Nutr Metab (Lond)* 9: 82.
10. Alleman RJ, Harvey IC, Farney TM, Bloomer RJ (2013) Both a traditional and modified Daniel Fast improve the cardio-metabolic profile in men and women. *Lipids Health Dis* 12: 114.
11. Bloomer RJ, Gunnels TA, Schriefer JM (2015) Comparison of a Restricted and Unrestricted Vegan Diet Plan with a Restricted Omnivorous Diet Plan on Health-Specific Measures. *Healthcare (Basel)* 3: 544-555.
12. Bloomer RJ, Kabir MM, Trepanowski JF, Canale RE, Farney TM (2011) A 21 day Daniel Fast improves selected biomarkers of antioxidant status and oxidative stress in men and women. *Nutr Metab (Lond)* 8: 17.
13. Bloomer RJ, Schriefer JHM, Gunnels TA, Lee SR, Sable HJ, et al. (2018) Nutrient Intake and Physical Exercise Significantly Impact Physical Performance, Body Composition, Blood Lipids, Oxidative Stress, and Inflammation in Male Rats. *Nutrients* 10.
14. Huang RY, Huang CC, Hu FB, Chavarro JE (2016) Vegetarian Diets and Weight Reduction: a Meta-Analysis of Randomized Controlled Trials. *J Gen Intern Med* 31: 109-116.
15. Yao M, Roberts SB (2001) Dietary energy density and weight regulation. *Nutr Rev* 59: 247-258.
16. Foright RM, Presby DM, Sherk VD, Kahn D, Checkley LA, et al. (2018) Is regular exercise an effective strategy for weight loss maintenance? *Physiol Behav* 188: 86-93.
17. Chastin SFM, De Craemer M, De Cocker K, Powell L, Van Cauwenberg J, et al. (2019) How does light-intensity physical activity associate with adult cardiometabolic health and mortality? Systematic review with meta-analysis of experimental and observational studies. *Br J Sports Med* 53: 370-376.
18. Swift DL, McGee JE, Earnest CP, Carlisle E, Nygard M, et al. (2018) The Effects of Exercise and Physical Activity on Weight Loss and Maintenance. *Prog Cardiovasc Dis* 61: 206-213.
19. Cao JJ (2018) Caloric restriction combined with exercise is effective in reducing adiposity and mitigating bone structural deterioration in obese rats. *Ann N Y Acad Sci* 1433: 41-52.
20. Kim MK, Cho SW, Park YK (2012) Long-term vegetarians have low oxidative stress, body fat, and cholesterol levels. *Nutr Res Pract* 6: 155-161.
21. Niemann B, Rohrbach S, Miller MR, Newby DE, Fuster V, et al. (2017) Oxidative Stress and Cardiovascular Risk: Obesity, Diabetes, Smoking, and Pollution: Part 3 of a 3-Part Series. *J Am Coll Cardiol* 70: 230-251.
22. Uttara B, Singh AV, Zamboni P, Mahajan R (2009) Oxidative stress and neurodegenerative diseases: a review of upstream and downstream antioxidant therapeutic options. *Curr Neuropharmacol* 7: 65-74.
23. Matsuda M, Shimomura I (2013) Increased oxidative stress in obesity: implications for metabolic syndrome, diabetes, hypertension, dyslipidemia, atherosclerosis, and cancer. *Obes Res Clin Pract* 7: 330-341.
24. Estadella D, Claudia da Penha Oller do MN, Oyama LM, Ribeiro EB, Damaso AR, et al. (2013) Lipotoxicity: effects of dietary saturated and trans fatty acids. *Mediators Inflamm* 2013: 137579.
25. Bonnefont-Rousselot D (2002) Glucose and reactive oxygen species. *Curr Opin Clin Nutr Metab Care* 5: 561-568.
26. Yadav RK, Singh M, Roy S, Ansari MN, Saeedan AS, et al. (2018) Modulation of oxidative stress response by flaxseed oil: Role of lipid peroxidation and underlying mechanisms. *Prostaglandins Other Lipid Mediat* 135: 21-26.
27. Diniz YS, Cicogna AC, Padovani CR, Silva MD, Faine LA, et al. (2003) Dietary restriction and fibre supplementation: oxidative stress and metabolic shifting for cardiac health. *Can J Physiol Pharmacol* 81: 1042-1048.
28. Yu Y, Gao Q, Xia W, Zhang L, Hu Z, et al. (2018) Association between Physical Exercise and Biomarkers of Oxidative Stress among Middle-Aged and Elderly Community Residents with Essential Hypertension in China. *Biomed Res Int* 2018: 4135104.



- Advances In Industrial Biotechnology | ISSN: 2639-5665
- Advances In Microbiology Research | ISSN: 2689-694X
- Archives Of Surgery And Surgical Education | ISSN: 2689-3126
- Archives Of Urology
- Archives Of Zoological Studies | ISSN: 2640-7779
- Current Trends Medical And Biological Engineering
- International Journal Of Case Reports And Therapeutic Studies | ISSN: 2689-310X
- Journal Of Addiction & Addictive Disorders | ISSN: 2578-7276
- Journal Of Agronomy & Agricultural Science | ISSN: 2689-8292
- Journal Of AIDS Clinical Research & STDs | ISSN: 2572-7370
- Journal Of Alcoholism Drug Abuse & Substance Dependence | ISSN: 2572-9594
- Journal Of Allergy Disorders & Therapy | ISSN: 2470-749X
- Journal Of Alternative Complementary & Integrative Medicine | ISSN: 2470-7562
- Journal Of Alzheimers & Neurodegenerative Diseases | ISSN: 2572-9608
- Journal Of Anesthesia & Clinical Care | ISSN: 2378-8879
- Journal Of Angiology & Vascular Surgery | ISSN: 2572-7397
- Journal Of Animal Research & Veterinary Science | ISSN: 2639-3751
- Journal Of Aquaculture & Fisheries | ISSN: 2576-5523
- Journal Of Atmospheric & Earth Sciences | ISSN: 2689-8780
- Journal Of Biotech Research & Biochemistry
- Journal Of Brain & Neuroscience Research
- Journal Of Cancer Biology & Treatment | ISSN: 2470-7546
- Journal Of Cardiology Study & Research | ISSN: 2640-768X
- Journal Of Cell Biology & Cell Metabolism | ISSN: 2381-1943
- Journal Of Clinical Dermatology & Therapy | ISSN: 2378-8771
- Journal Of Clinical Immunology & Immunotherapy | ISSN: 2378-8844
- Journal Of Clinical Studies & Medical Case Reports | ISSN: 2378-8801
- Journal Of Community Medicine & Public Health Care | ISSN: 2381-1978
- Journal Of Cytology & Tissue Biology | ISSN: 2378-9107
- Journal Of Dairy Research & Technology | ISSN: 2688-9315
- Journal Of Dentistry Oral Health & Cosmesis | ISSN: 2473-6783
- Journal Of Diabetes & Metabolic Disorders | ISSN: 2381-201X
- Journal Of Emergency Medicine Trauma & Surgical Care | ISSN: 2378-8798
- Journal Of Environmental Science Current Research | ISSN: 2643-5020
- Journal Of Food Science & Nutrition | ISSN: 2470-1076
- Journal Of Forensic Legal & Investigative Sciences | ISSN: 2473-733X
- Journal Of Gastroenterology & Hepatology Research | ISSN: 2574-2566
- Journal Of Genetics & Genomic Sciences | ISSN: 2574-2485
- Journal Of Gerontology & Geriatric Medicine | ISSN: 2381-8662
- Journal Of Hematology Blood Transfusion & Disorders | ISSN: 2572-2999
- Journal Of Hospice & Palliative Medical Care
- Journal Of Human Endocrinology | ISSN: 2572-9640
- Journal Of Infectious & Non Infectious Diseases | ISSN: 2381-8654
- Journal Of Internal Medicine & Primary Healthcare | ISSN: 2574-2493
- Journal Of Light & Laser Current Trends
- Journal Of Medicine Study & Research | ISSN: 2639-5657
- Journal Of Modern Chemical Sciences
- Journal Of Nanotechnology Nanomedicine & Nanobiotechnology | ISSN: 2381-2044
- Journal Of Neonatology & Clinical Pediatrics | ISSN: 2378-878X
- Journal Of Nephrology & Renal Therapy | ISSN: 2473-7313
- Journal Of Non Invasive Vascular Investigation | ISSN: 2572-7400
- Journal Of Nuclear Medicine Radiology & Radiation Therapy | ISSN: 2572-7419
- Journal Of Obesity & Weight Loss | ISSN: 2473-7372
- Journal Of Ophthalmology & Clinical Research | ISSN: 2378-8887
- Journal Of Orthopedic Research & Physiotherapy | ISSN: 2381-2052
- Journal Of Otolaryngology Head & Neck Surgery | ISSN: 2573-010X
- Journal Of Pathology Clinical & Medical Research
- Journal Of Pharmacology Pharmaceutics & Pharmacovigilance | ISSN: 2639-5649
- Journal Of Physical Medicine Rehabilitation & Disabilities | ISSN: 2381-8670
- Journal Of Plant Science Current Research | ISSN: 2639-3743
- Journal Of Practical & Professional Nursing | ISSN: 2639-5681
- Journal Of Protein Research & Bioinformatics
- Journal Of Psychiatry Depression & Anxiety | ISSN: 2573-0150
- Journal Of Pulmonary Medicine & Respiratory Research | ISSN: 2573-0177
- Journal Of Reproductive Medicine Gynaecology & Obstetrics | ISSN: 2574-2574
- Journal Of Stem Cells Research Development & Therapy | ISSN: 2381-2060
- Journal Of Surgery Current Trends & Innovations | ISSN: 2578-7284
- Journal Of Toxicology Current Research | ISSN: 2639-3735
- Journal Of Translational Science And Research
- Journal Of Vaccines Research & Vaccination | ISSN: 2573-0193
- Journal Of Virology & Antivirals
- Sports Medicine And Injury Care Journal | ISSN: 2689-8829
- Trends In Anatomy & Physiology | ISSN: 2640-7752

Submit Your Manuscript: <https://www.heraldopenaccess.us/submit-manuscript>